

Mathematical Preliminaries

Sets:

- An unordered collection of elements (order doesn't matter).
- Can be finite, $\{2, 3, 4\}$, or infinite $\{1, 2, 3, 4, \dots\}$.
- Set **membership**: \in, \notin

Ex: $4 \in \{2, 3, 4\}$, $1 \notin \{2, 3, 4\}$,

- Sets can contain other sets: $\{2, \{5\}\}$, $\{\{0\}\} \neq \{0\} \neq 0$
- Two sets are equal if they contain the same elements.

Common Sets

- Naturals: $N = \{0, 1, 2, 3, 4, \dots\}$
- Integers: $Z = \{\dots - 2, -1, 0, 1, 2, \dots\}$
- Rationals: $Q = \{\frac{a}{b} \mid a, b \in Z, b \neq 0\}$
- Reals: R
- Empty set: $\emptyset = \{\}$
- Set **definition**: ' \mid ' means “such that”.

Ex: $\{k \mid k \in N, 0 < k < 4\}$

Set operations

- Subset: \subseteq, \subset .
- $\forall S, \emptyset \subseteq S$.
- $\forall S, S \subseteq S$.
- Union (\cup), Intersection (\cap).
- **Set difference:** $S - T = \{x | x \in S \wedge x \notin T\}$.
- **Set complement:** $\neg S$ or $\bar{S} = \{x | x \notin S\} = U - S$, where U is a universal set (everything).
- **Disjoint** sets: $S \cap T = \emptyset$.

Set Cardinality

- **Cardinality:** $|S|$ = number of elements in S .
- **Power set** of a set A , 2^A is the set of all subsets of A .

Example: $A = \{2, 3\}$, then the power set of A is $2^A = \{\emptyset, \{2\}, \{3\}, \{2, 3\}\}$.

Question: if $|A| = n$, what is the cardinality of the power set? Answer: 2^n .

- **DeMorgan's laws:**

$$\neg(B \cap C) = \neg B \cup \neg C$$

$$\neg(B \cup C) = \neg B \cap \neg C$$

Cartesian product

- Given two sets A and B , the **Cartesian product** or **cross product** $A \times B$ is the set of all *ordered pairs* wherein the first element is a member of A and the second element is a member of B .

Example: if $A = \{1, 2\}$ and $B = \{x, y, z\}$, then
 $A \times B = \{(1, x), (1, y), (1, z), (2, x), (2, y), (2, z)\}$.

Question: what is the cardinality of $A \times B$? Answer:
 $|A| \times |B|$.

Binary relations

- A **binary relation** R on two sets A and B is a subset of the Cartesian product $A \times B$. If $(a, b) \in R$, this is equivalently written as aRb .
- Types of relations $R \subseteq A \times A$:
 - *reflexive*: aRa , for all $a \in A$
 - *symmetric*: $aRb \Rightarrow bRa$, for all $a, b \in A$
 - *transitive*: aRb and $bRc \Rightarrow aRc$, for all $a, b, c \in A$
 - *equivalence*: reflexive and symmetric and transitive.
- Examples: $<$, \geq , $=$.

Functions

- A **function** $f : A \rightarrow B$ is a binary relation on A and B such that for all $a \in A$, there is one and only one $b \in B$ such that $(a, b) \in f$.

$(a, b) \in f$ is equivalently written $f(a) = b$.

A is called f 's **domain** and B is the **codomain**.

We say that a is the **argument** of f and that $f(a) = b$ is the **value (image)** of f at a .

- The **range** of f is the image of its domain, that is, $f(A) = \{b \in B : b = f(a) \text{ for some } a \in A\}$.
- A function is a **surjection** if its range is its codomain.

Functions (cont'd)

- A function $f : A \rightarrow B$ is an **injection (one-to-one)** if distinct arguments to f produce distinct values, that is, if $a \neq a'$ implies $f(a) \neq f(a')$.

Example:

- A function $f : A \rightarrow B$ is a **bijection (one-to-one correspondence)** if it is injective and surjective.

Example:

Floor, Ceiling

floor and ceiling:

- Let $x \in R$, then:
 - $\lfloor x \rfloor =$ largest integer $\leq x$ — “floor”. (e.g., $\lfloor 8.2 \rfloor = 8$)
 - $\lceil x \rceil =$ smallest integer $\geq x$ — “ceiling”. (e.g., $\lceil 8.2 \rceil = 9$)

Basic facts:

- $x - 1 < \lfloor x \rfloor \leq x \leq \lceil x \rceil < x + 1$
- If n is a integer then $\lfloor n/2 \rfloor + \lceil n/2 \rceil = n$

$$\lceil \frac{\lfloor \frac{n}{2} \rfloor}{2} \rceil = \lceil \frac{n}{4} \rceil$$

Polynomial and Exponential

Polynomials:

$$p(n) = \sum_{k=0}^d a_k \cdot n^k = a_d \cdot n^d + \dots a_1 \cdot n + a_0 \quad (1)$$

Exponential Function:

$$a^0 = 1$$

$$a^1 = a$$

$$a^{-1} = 1/a$$

$$(a^m)^n = a^{mn}$$

$$a^m a^n = a^{m+n}$$

Logarithms

Logarithms:

- definitions: $\lg n = \log_2 n$, $\ln n = \log_e n$
- $\log_c ab = \log_c a + \log_c b$.
- $\log_c a^b = b \cdot \log_c a$.
- $\log_c \frac{a}{b} = \log_c a - \log_c b$.
- $\log_c a = \frac{\log_d a}{\log_d c}$. (change base)
- $a^{\log_c n} = n^{\log_c a}$
- derivatives: $(\ln a)' = \frac{1}{a}$, $(\lg a)' = \frac{\lg e}{a}$.

Factorial

Factorials:

$$n! = \begin{cases} 1 & \text{for } n = 0 \\ n(n-1)! & \text{for } n > 0 \end{cases}$$

Note:

- $n! \leq n^n$
- $\sqrt{2\pi n} \cdot \left(\frac{n}{e}\right)^n \leq n! \leq \sqrt{2\pi n} \cdot \left(\frac{n}{e}\right)^{n+\left(\frac{1}{12}n\right)}$

The last formula is called “Stirling’s approximation” for $n!$.

Summation & Recurrences

Summations

Given a sequence of numbers $a_1, a_2, a_3, \dots, a_n$, the summation $a_1 + a_2 + \dots + a_n$ is written as

$$\sum_{i=1}^n a_i$$

The infinite sum $a_1 + a_2 + \dots$ is written as

$$\sum_{i=1}^{\infty} a_i$$

and it is formally interpreted as

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n a_i.$$

General Properties of Summations

Linearity

$$\sum_{k=1}^n (ca_k + b_k) = c \sum_{k=1}^n a_k + \sum_{k=1}^n b_k.$$

Arithmetic Series

$$\sum_{k=1}^n k = \frac{1}{2}n(n+1) = \Theta(n^2).$$

Sum of squares

$$\sum_{k=1}^n k^2 = \frac{1}{6}n(n+1)(2n+1) = \Theta(n^3).$$

Series

Sum of cubes

$$\sum_{k=1}^n k^3 = \frac{n^2(n+1)^2}{4} = \Theta(n^4).$$

Geometric Series For real number $x \neq 1$,

$$\sum_{k=0}^n x^k = 1 + x + x^2 + x^3 + \dots + x^n = \frac{x^{n+1} - 1}{x - 1}.$$

The following geometric series are used frequently:

$$\sum_{k=0}^n 2^k = \frac{2^{n+1} - 1}{2 - 1} = 2^{n+1} - 1.$$

$$\sum_{k=0}^{\infty} x^k = \frac{1}{1 - x} \quad (\text{if } |x| < 1)$$

More Series

Using integrals:

- if f is a continuous, increasing function:

$$\int_{a-1}^b f(x)dx \leq \sum_{i=a}^b f(i) \leq \int_a^{b+1} f(x)dx$$

- if f is a continuous, decreasing function:

$$\int_a^{b+1} f(x)dx \leq \sum_{i=a}^b f(i) \leq \int_{a-1}^b f(x)dx$$

- Example: $f(k) = \frac{1}{k}$

$$\ln(n+1) \leq \sum_{i=1}^n \frac{1}{k} \leq \ln(n) + 1, \quad \sum_{i=1}^n \frac{1}{k} = \ln(n) + O(1),$$

Graphs

- A **directed graph** G is a pair (V, E) , where V is the set of vertices, and E is the set of edges (i.e. ordered pairs of vertices).


Review: *adjacency, in-degree, out-degree, path, cycle.*

- In an **undirected graph** $G = (V, E)$, the edges are unordered pairs of vertices.

Review: *adjacency, degree, path, cycle.*

Review on Graphs

$G=(V,E)$



$$V = \{1, 2, 3, 4, 5\}$$

$$E = \{(1, 2), (2, 3), (3, 5), (5, 4), (4, 1)\}$$

Representation of graphs

- Adjacency List
- Adjacency Matrix

Review on Trees

- A **free tree** is a connected, acyclic, undirected graph.
- A **rooted tree** is a free tree in which one vertex (the root) is distinguished from the others.

Review: *ancestor/descendant, parent/child, siblings, external/internal nodes, depth & height.*

Proofs

Mathematical Statements:

- Definition, Lemma, Theorem, Corollary

Types of Proofs:

- Contradiction
- Induction
- Counter-example

Proof by Contradiction

Example: $\sqrt{2}$ is rational.

Proof by Induction

If we want to prove a statement $P(n)$ is true for all natural numbers $n \in \{1, 2, 3, \dots\}$, we can achieve this with the following two steps:

- 1 Prove that the statement holds when $n = 1$
($P(1)$ is true). — — — **basis**
- 2 Prove that if the statement holds for $n = m$, then the same statement holds for $n = m + 1$.
($P(m) \Rightarrow P(m + 1)$). — — — **induction step**

Example

$$\sum_{1}^n = \frac{n(n+1)}{2} \text{ for } n = \{1, 2, 3\dots\}.$$

Proof by Induction: Generalizations

Generalization type 1:

- If we want to prove a statement P not for all natural numbers but only for all numbers greater than a certain number b then the following two steps are sufficient
 1. **basis:** Prove that the statement holds when $n = b$.
 2. **induction step:** Prove that if the statement holds for $n = m$ then the same statement also holds for $n = m + 1$.

Generalizations

Generalization type 2:

- Another generalization allows that in the second step, we not only assume that the statement holds for $n = m$ but also for all n smaller than or equal to m . This leads to the following two steps.
 1. **basis:** Prove that the statement holds when $n = b$.
 2. **induction step:** Prove that if the statement holds for $n \leq m$ then the same statement also holds for $n = m + 1$.

Example

Every natural number greater than 1 is a product of prime numbers.